


This article was downloaded by: [University of Nebraska, Lincoln]

On: 06 April 2015, At: 10:28

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Critical Reviews in Food Science and Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/bfsn20>

### WHEAT ALLERGY & INTOLERANCE; RECENT UPDATES AND PERSPECTIVES

Imran Pasha<sup>a</sup>, Farhan Saeed<sup>a</sup>, Muhammad Tauseef Sultan<sup>b</sup>, Rizwana Batool<sup>a</sup>, Mahwash Aziz<sup>a</sup> & Waqas Ahmed<sup>a</sup>

<sup>a</sup> National Institute of Food Science and Technology, University of Agriculture, Faisalabad

<sup>b</sup> Department of Food Sciences, Bahauddin Zakariya University, Multan

Accepted author version posted online: 02 Sep 2013.



CrossMark

[Click for updates](#)

To cite this article: Imran Pasha, Farhan Saeed, Muhammad Tauseef Sultan, Rizwana Batool, Mahwash Aziz & Waqas Ahmed (2013): WHEAT ALLERGY & INTOLERANCE; RECENT UPDATES AND PERSPECTIVES, Critical Reviews in Food Science and Nutrition, DOI: [10.1080/10408398.2012.659818](https://doi.org/10.1080/10408398.2012.659818)

To link to this article: <http://dx.doi.org/10.1080/10408398.2012.659818>

Disclaimer: This is a version of an unedited manuscript that has been accepted for publication. As a service to authors and researchers we are providing this version of the accepted manuscript (AM). Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). During production and pre-press, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

# ACCEPTED MANUSCRIPT

**TYPE OF ARTICLE:** Review article

**TITLE:** WHEAT ALLERGY & INTOLERANCE; RECENT  
UPDATES AND PERSPECTIVES

**1. Imran Pasha**

*National Institute of Food Science and Technology,  
University of Agriculture, Faisalabad*

**2. Farhan Saeed**

*National Institute of Food Science and Technology,  
University of Agriculture, Faisalabad*

**3. Muhammad Tauseef Sultan**

*Department of Food Sciences  
Bahauddin Zakariya University, Multan*

**4. Rizwana Batool**

*National Institute of Food Science and Technology,  
University of Agriculture, Faisalabad*

**5. Mahwash Aziz**

*National Institute of Food Science and Technology,  
University of Agriculture, Faisalabad*

**6. Waqas Ahmed**

*National Institute of Food Science and Technology,  
University of Agriculture, Faisalabad*

# ACCEPTED MANUSCRIPT

†Corresponding author

**Rizwana Batool**

*National Institute of Food Science and Technology,*

*University of Agriculture,*

*Faisalabad*

Contact No. 0092-333-6724514

Email: [rizwana\\_uaf@yahoo.com](mailto:rizwana_uaf@yahoo.com)

WHEAT ALLERGY & INTOLERANCE; RECENT UPDATES AND PERSPECTIVES

## Abstract

The current review article highlights the complications associated with communities relying on wheat as their dietary staple. Although, wheat is an important source of nutrients but is also linked with allergenic responses in genetically susceptible subjects. The wheat proteins especially  $\alpha$ -amylase inhibitors,  $\omega$ -5 gliadins, prolamins, non-prolamin, glucoprotein and profilins are of significance importance. The allergenic responses are further categorized into IgE-mediated and non-IgE-mediated reactions. Conjugation and degranulation of the IgEs with the allergens results in release of several mediators. In contrary, non-IgE-mediated wheat allergy depends on immune complexes formed by food and food antibodies and cell-mediated immunity. As results, different diseases tend to occur on the completion of these reactions i.e. celiac disease (CD), baker's asthma, diarrhea, atopic dermatitis (AD) and urticaria. This instant article highlighted the concept of food allergy with special reference to wheat. The models are developed that are included in this article showing the wheat allergen, their possible routes, impact on human health, and indeed possible remedies. The article would provide the basic information for the researchers, common man, and allied stakeholders to cater the issue in details. However, the issue needs the attention of the researchers as there is a need to clarify the issues of wheat allergy and wheat intolerance.

**Keywords:** Wheat allergens, Immune cells, Celiac disease, Baker's asthma, Atopic dermatitis

## Introduction

Provision of adequate and balanced food is important for vitality of life. However, adverse reactions on foods are extremely common and some of them are described as allergic that imply an immune mechanism (Guandalini and Newland, 2011). In this regard, Food allergy is defined as an undesirable response of the immune system against proteins. Generally, foods that induce rhinitis, sinusitis, bronchitis, asthma, intestinal cramps, diarrhea, hives, angioedema, eczema and migraines are said to be "allergenic". There have been increasing reports of food allergy to a wide range of foods but cow's milk, egg, peanut, wheat, soy, tree nuts, fish and shell-fish are of critical significance (Sampson, 2004). These foods generally account for nearly 90% of the food-allergic reactions. However, the prevalence of apparent food hypersensitivity varies (1.4–19.1%) among different communities. These wide variations can also be attributed to differences in the diagnostic criteria, study design and population etc. (Crespo and Rodriguez, 2003). In general, about 1.5% of adults and 5 to 8% of growing children suffer from food allergies (Sampson, 2000). The estimated percentage of allergies caused by different foods is 0 to 3% for milk, up to 1.7% for egg, 0.2 to 1.6% for peanut and 1 to 10.8% for other foods (Zuidmeer et al., 2008). The allergies due to consumption of cereals are another major problem in many countries where they cereals are habitually consumed (Battais et al., 2008). In Japan, wheat is third commonest food causing allergy in children following eggs and cow` s milk (Imai and Iikura, 2003). Though allergies to

cow's milk and egg can be tolerated by elimination, wheat allergy can't be controlled through elimination. In Pakistan, self-reported frequency of allergy varied from 1.2 to 17% for milk, 0.2 to 7% for egg, 0 to 2% for peanut and fish, 0 to 10% for shellfish and 3 to 35% for other foods (Rona et al., 2007).

Food allergy is also dependent upon genetic factors, cultural and dietary habits and exposure to allergenic products. For instance, peanut allergy is common in UK, France and Switzerland but rare in Italy, Singapore and Israel (Dalal et al., 2002). Food allergies have augmented considerably in the past decade. At the onset, food intolerance must be separated from food allergies and, furthermore, these allergies should be categorized into an IgE, Non-IgE, or a mixed response. The clinical features differ from life-threatening anaphylaxis to milder IgE-mediated reactions, atopic dermatitis, and gastrointestinal symptoms (Heratizadeh et al., 2011; Wang et al., 2011). As far as IgE-mediated wheat allergy is concerned, specific IgE are developed in the absence of oral tolerance (OT). IgE are related to high affinity specific Fc receptors FcεRI present on the mast cells, fixed cells present in the mucosa and skin, and basophils circulating in the blood (Pizzuti et al., 2011). These cells have various cytoplasmic granules mainly including histamine in which preformed mediators are stored. When the allergenic proteins are captivated in the gut they contact with specific IgEs linked to mast cells/basophils. The conjugation of the IgEs with the allergens triggers these cells to degranulate, thus releasing immuno-mediators in the surrounding

microenvironment (Sotkovsky et al., 2011). Consequently, different mediators such as prostaglandins, leukotriens, and cytokines are also released. An abrupt reaction follows few minutes after making contact with the allergen essentially due to the histamine release. At the base of the reaction there is vasodilation, tissue fluid exudation, smooth muscle contraction and mucous secretion. A late-phase response follows the instant reaction which begins 4–6 hours after contact with the allergen and continues for several days. This response is caused by chemotactic mediators released at the same time as the immediate reaction which promote selective conscription of inflammatory cells, mainly eosinophils and neutrophils. These cells infiltrate the tissues produce inflammation lasting for a period of four to five days. However, in order to predict IgE mediated allergy, the two clinical elements are required i.e. presence of IgE specific antibodies and a proven relationship between ingestion of the food and the appearance of the symptoms (Cabrera-Chavez et al., 2009).

On the other hand, non-IgE-mediated food allergy consists of immune reactions that are dependent upon presence of antibodies other than IgE (e.g. IgG, IgM and IgA) and immune complexes (food, food antibodies, and cell-mediated immunity). It is important to remember that there is no verification to support the affiliation between the allergenic food and a reaction to non-IgE antibodies in any allergic disease. However, in normal individuals, there is an increase in IgG food antibodies after

ingestion of the food. Likewise, there is no solid evidence about the responsibility of cell-mediated immunity (Paganelli et al., 1987).

Strategies for the prevention of allergy have been proposed including the utilization of product with extensively reduced allergenicity. A perfect history is vital in approaching the administration (Wantanabe, 1993). Structural proteins are imperative determinants of the sternness of the reactions and may often foresee the natural history and cross reactivity. Indicative work up must be directed by the clinical history. Skin testing and food-specific IgE done by standard methods are very helpful, while oral tolerance (OT) may be specified in some circumstances (Ramesh, 2008).

## **Wheat allergy**

Wheat is widely consumed cereal all over the world and its potential to cause allergenic responses is a matter of great interest for clinical practitioners and nutritionists. Despite the large consumption of wheat all over the world, there are few reports about clinical aspects of wheat food allergy in controlled adults. Wheat is amongst the most ordinary reasons of food-related allergies and intolerances that are linked with its ingestion or inhalation (Yacoub et al., 2011). Wheat, like all other foods, contains a number of proteins (more than 100), of which, some have been identified as allergens. The type and extent of proteins in a cereal have a major impact in determining the quality and end use properties of the cereal (Shewry and Halford, 2002). Wheat food allergy is

frequent in children and infants and indeed is listed as one of the six 6 most commonly implicated food allergens (FAO, 1995). The wheat allergy has been confirmed by the large number of double-blind, placebo-controlled food challenges (DBPCFCs) performed in subjects over the globe (Palosuo et al., 2001).

On the basis of solubility, wheat proteins can be characterized as water-soluble, salt-soluble, alcohol-soluble and alcohol-insoluble (Pomeranz, 1988). According to the type of wheat, proteins (albumin, globulin and gluten) vary in proportion and this variability results in non-consistent reactions in different wheat products (Joneja, 1998). Among gluten protein, gliadins (28-42%), the major prolamin protein in wheat is soluble in 70-90% alcohol whereas glutenins (42-62.5%), the major glutelin proteins in wheat is soluble in dilute acid or alkali solutions (Armentia et al., 2002). Wheat proteins are subdivided into the water/salt-soluble (including albumins and globulins) and insoluble (containing gliadins and glutenins) fractions. The latter represent about 80% of all wheat proteins. Mono-meric gliadins mainly exhibit intramolecular disulfide linkages, thus being compressed and of round shape. The different gliadin subunits share amino acid sequence and 3-dimensional structure homologies (Bietz, 1977). Several wheat salt-soluble proteins (albumins and globulins), alpha-amylase/trypsin inhibitors, peroxidase, thioredoxin, nonspecific lipid transfer protein, serine proteinase inhibitor, and thaumatin-like protein-as well as salt-insoluble storage proteins (prolamins, namely, gliadins and glutenins) act as allergens (Salcedo et al., 2011).

Recently, thaumatin-like protein (TLP), and lipid transfer protein 2G are also responsible for allergenic responses resulting due to wheat ingestion. Gluten proteins especially its gliadin fraction is the major aspect in triggering celiac disease, a common enteropathy, linked with intake of wheat gluten proteins and related prolamins from oat, rye and barley (Mamone et al., 2011). Gliadin is known as a presumptive major allergen by immunoblotting and RAST inhibition test (Morita et al., 2001). Previously, Palosuo et al. (2001) have identified  $\omega$ -5 gliadin, which had been reported g-gliadin-like protein in their preceding report (Palosuo et al., 1999) as a major allergen in children with wheat-dependent exercise induced anaphylaxis (Palosuo et al., 2001). Wheat proteins belonging to both the soluble and insoluble fractions can act as allergens in sensitized individuals. In this regard, the soluble proteins seem to be responsible for respiratory allergy, such as baker's asthma arising from flour inhalation.

By characterization, gluten is found only in wheat, although the term is commonly used to refer to any similar prolamins protein in any grain (Thompson, 2000). The percentages of different types of prolamins found in various cereals have been given in Table 1. Because the type and proportion of prolamins proteins in these grains vary, the kind of reaction (if any) they are expected to cause also varies (Fasano and Catassi, 2001). An elevated IgE in gliadin-intolerant patients is indicated in some studies (Varjonen *et al.*, 2000), however, little is known about cellular mechanisms involved in such conditions. The generation of specific IgE against gliadin in atopic patients suggested that IgE

dysregulation is related to gliadin intolerance (Giannotti et al., 2001). The IgE production by human  $\beta$ -lymphocytes is mainly controlled by interleukin (IL)-4 (De Vries et al., 1991) produced by TH2-lymphocytes, mast cells or basophils (Arock et al., 1993) and by CD40-CD40 ligand interactions between  $\beta$  and T-lymphocytes (Vercelli, 1995). IL-4 is up-regulated during gliadin intolerance (Troncone et al., 1998) that further modulates the phenotype and function of  $\beta$ - and T-lymphocytes and monocytes/macrophages (Dugas et al., 1990). In  $\beta$  cells and monocytes/macrophages, IL-4 induces CD23-expression which amplifies IL-4-induced IgE production. In addition to these immunological mechanisms, it is shown that cellular redox status also controls this production (Benngtsson et al., 2001). Indeed, the redox metabolism is responsible for controlling dendritic cell functions (Alderman et al., 2002), and thus it mediates the protective immunity (Bachmann and Kopf, 2002). Gliadin can induce a redox imbalance in gastro-intestinal epithelial cells and macrophages probably due to over-production of free radicals. According to a study, it was evaluated that native gliadin and gliadin lysates (GliIys) showed the regulatory effects on the spontaneous and the IL-4-induced IgE production by normal human peripheral blood mononuclear cells (PBMC). Although, enzymatic (Cu/Zn-superoxide dismutase) and non-enzymatic (thiols) antioxidants can perform pharmacological roles under such conditions (Ribavene et al., 1999) but imbalance in redox potential results in severity of allergenic responses. IgE-mediated hypersensitivity reactions for ingested foods have been an important health problem. Several hypersensitivity reactions have been reported that are concerned with

orally ingested wheat (Sampson, 1983). It is hypothesized that IgE from atopic dermatitis (AD) patients react with wheat and other cereal allergens. Number of patients suspected of AD owing to wheat ingestions is increasing in Japan. Multiple findings recommended that some wheat proteins (salt-soluble and salt insoluble fractions) are reactive to IgE in AD patients (Miyakawa et al., 1988).

Generally, food allergic reactions to wheat can give way to an array of clinical manifestations that can range from immediate to delayed, and their strictness can vary from mild to life-threatening. Typical immediate symptoms include erythema, pruritus, eczema, gastrointestinal reactions, oropharyngeal symptoms, urticaria, angioedema, atopic dermatitis, rhinitis, asthma, and anaphylaxis (Heffler et al., 2011; Bock, 1986). Investigators have confirmed that IgE antibodies specific to wheat albumins and globulins are present in the serum from children with wheat allergy and patients with baker's asthma (Cho et al., 2011; Jones et al., 1995).

In infants and young children, the symptoms associated with wheat allergy usually arise after weaning. In children, gastrointestinal problems are more evident including abnormal stools, abdominal distension & pain, flatulence, nausea, vomiting and intestinal malabsorption (Alderman et al., 2002). Children are also suffered by irritability, apathy, loss of appetite, weight loss, poor weight gain, short stature, muscle wasting, hypotonia, general failure to thrive, poor school performance, bone and joint pains, and occasionally rickets (Varjonen et al., 2000).

In older children and adults, the symptoms may quite vary ranging from severe weight loss, diarrhoea and bulky, offensive stools to less severe symptoms. Subtle complaints of abdominal bloating, cramping, flatulence and constipation are often mistakenly attributed to irritable bowel syndrome. Some individuals suffer from only anemia-related fatigue and have no gastrointestinal symptoms. Other manifestations of the disease include osteopenic bone disease, infertility, tetany, ataxia and neurologic disorders. Recent studies show the presence of ataxia in persons with serological evidence of gluten sensitivity but without overt gastrointestinal symptoms or evidence of small-bowel inflammation (Joneja, 1998). The details of wheat allergens and various allergenic disorders are discussed in detail herein.

**Place figure 1 here**

## **WHEAT ALLERGENS**

Several wheat proteins have been described as allergens on the molecular level. Amongst,  $\alpha$ -amylase inhibitors, acyl-coenzyme, oxidase, fructose-biphosphate aldolase, and wheat-flour peroxidase &  $\omega$ -5 gliadin are of significance importance.

### **$\alpha$ -amylase inhibitors**

Several groups of investigators have reported that some of the wheat allergens responsible for baker's asthma are members of a family of wheat  $\alpha$ -amylase inhibitors

(Sandiford et al., 1995). A 16 kDa wheat protein, a glycosylated subunit of the wheat tetrameric  $\alpha$ -amylase inhibitor, has been identified and confirmed as an important allergen. Using IgE immune-blotting, Pfeil et al. (1990) identified major  $\alpha$ -amylase inhibitor with molecular weights of 15 and 17 kDa. In similar investigations, Sandiford et al. (1995) identified 15 and 12 kDa  $\alpha$ -amylase inhibitor in patients with baker's asthma. In another study, Jones et al. (1995) showed 20 and 47 kDa wheat proteins as potential allergens in children with confirmed wheat hypersensitivity. Ikezawa et al. (1994) found 33-98 kDa IgE-binding wheat proteins with AD patients sera in the salt-insoluble fraction by immunoblotting. In contrast, Ikezawa et al. (1994) reported that salt-soluble proteins with molecular masses of about 14 kDa bound IgE in AD patients sera, and these might be the same proteins considered to be major allergens of baker's asthma. Likewise, James et al. (1997) identified a subunit of the wheat dimeric  $\alpha$ -amylase inhibitor which binds IgE in patients with hypersensitivity reactions to ingested wheat proteins. In another study, Gomez et al. (1990) revealed that CM3 (14 kd), one of the subunits of  $\alpha$ -amylase inhibitors, binds to IgE sera from AD patients. The single allergen sensitization profiles obtained with 17 recombinant wheat flour allergens while two CCDs revealed no major allergen for German bakers. The highest frequencies were found for  $\alpha$ -amylase inhibitors and CCDs (Sander et al., 2011)

Walsh and Howden, (1989) using an epitope mapping method, have recently identified a putative allergenic peptide in the NH<sub>2</sub>-terminal amino acid sequence

of protein (0.28). It was further considered as a new wheat monomeric inhibitor of heterologous  $\alpha$ -amylases. Similarly, a major allergen with activity against insect  $\alpha$ -amylase has been also characterized in barley flour (Tordesillas et al., 2011; Barber et al., 1989). Both aforementioned proteins have around 14 kDa, and belong to the same protein family in cereals, which includes trypsin inhibitors and subunits of monomeric, dimeric and tetrameric inhibitors of heterologous  $\alpha$ -amylases.

## **Prolamins**

Prolamins, the second major type of protein are alcohol-soluble and water-insoluble storage proteins of cereal grains (Shewry, 2002). In addition, they often have unusual amino acid compositions, being particularly rich in proline and glutamine (Shewry and Tatham 1999). The prolamins themselves are immensely diverse in structure, with the major prolamins of maize ( $\alpha$ -zeins) and closely related panicoid' cereals (millets, sorghum) forming an apparently distinct group. In addition, wheat prolamins have also been implicated in dietary allergy, with symptoms including atopic dermatitis and exercise-induced anaphylaxis (Varjonen et al., 1997). Consequently; the prolamin currently forms the largest and most widely distributed group of plant food allergens.

## $\omega$ -5 Gliadins

The function of gliadin and its derived peptides in educing the unfavorable reactions in celiac disease is still far from being entirely elucidated. Gliadins proteins are further divided into subunits i.e.  $\alpha$ -,  $\beta$ -,  $\gamma$ - and  $\omega$ -gliadins. Whilst,  $\omega$ -gliadins are further subdivided into  $\omega$ 1,  $\omega$ 2 and  $\omega$ 5 components on the basis of sequences and mobility (Kasarda et al., 1983). Amongst,  $\alpha$ -,  $\beta$ -, and  $\gamma$ -gliadins are toxic to persons with ceolic disease (CD) and  $\omega$ -gliadins is non-toxic to persons with CD (Pomeranz, 1988).

Wheat  $\omega$ -5 gliadin has been identified as the major allergen in wheat-dependent, exercise-induced anaphylaxis (WDEIA) (Staden et al., 2007). More lately, it has been reported that IgE to  $\omega$ -5 gliadin is also highly predictive of immediate allergy to ingested wheat in children, with increased levels correlating with positive oral wheat challenge results (Bauer *et al.*, 1999). More than 80% of children with immediate symptoms to wheat reportedly had IgE anti-bodies to  $\omega$ -5 gliadin. The findings of Lauriere et al. (2006) and Patriarca et al. (2006) showed that sera from patients allergic to hydrolyzed wheat proteins with immediate skin reactions and from patients with WDEIA exhibited distinct patterns of reactivity. Similarly, Niggemann et al. (2006) have shown that different profiles of wheat antigens are known by patients with IgE-mediated wheat allergy and celiac disease (Tanaka et al., 2011; Palosuo et al., 2001). Amongst, w5-gliadins induced release of histamine from the basophils of patients with WDEIA but not from those of controls (Lehto et al., 2003).

Fast  $\omega$ -gliadin (contains  $\omega$ -5 gliadin) is a major allergen in wheat-dependent exercise-induced anaphylaxis and IgE against fast  $\omega$ -gliadin cross-reacts to other gliadins.  $\omega$ 5-gliadin has been identified as a major allergen in adults having WDEIA (Matsuo et al., 2004; Morita et al., 2003; Palosuo et al., 1999).

## **Non-Prolamin**

Non-prolamins consist of water-soluble albumins and salt-soluble globulins (Singh and Skerritt, 2001). Few studies on non-prolamins have been carried out so far as compared to gliadins and glutenins. Glutenins are composed of high molecular weight (HMW) subunits and low molecular weight (LMW) subunits. In fact, non-prolamins possess multiple functions during growth and development of wheat. For instance, albumins and globulins include enzymes and inhibitors of enzymes that regulate development at different stages. The relative amounts of essential amino acids such as aspartate, threonine, lysine and tryptophan for humans are more abundant in albumins and globulins, but less adequate than in storage proteins.

## **Glycoproteins**

Glycoproteins are proteins that contain oligosaccharide chains (glycans) covalently attached to polypeptide side-chains. The carbohydrate is attached to the protein in a co-translational or posttranslational modification that is known as glycosylation. In proteins that have segments extending extracellularly, the extracellular segments are

often glycosylated. Glycoproteins are often important integral membrane proteins, where they play a role in cell-cell interactions. Glycoproteins also occur in the cytosol, but their functions and the pathways producing these modifications in this compartment are less well-understood (Funakoshi and Suzuki, 2009). The major food allergens identified as class 1 allergens are water-soluble glycoproteins that are 10 to 70 kDa in size and fairly stable to heat, acid and proteases (Sampson, 1999). According to Robert et al. (2006), monosaccharides commonly found in eukaryotic glycoproteins are given in Table 2.

## **Profilins**

Profilins play a major role in the regulation of polymerization of actin filaments as well as consist of a large portion of the class 2 allergens and frequently show cross-reactivity between pollen and food (Breiteneder and Ebner, 2000). Patients often become sensitized to the inhaled pollen and, because of the cross-reactivity with profilins in the fruit or vegetable, experience oral and pharyngeal symptoms when ingesting the raw fruit or vegetable (i.e., the pollen-food allergy or oral allergy syndrome). The majority of these class 2 allergens are presumably comprised of conformational epitopes and therefore highly heat labile, susceptible to enzymatic degradation and difficult to isolate, often giving unsatisfactory standardized extracts for diagnostic purposes.

## **Derivative allergies**

Proteins are made of a chain of dehydrated amino acids. When enzymes cut proteins into pieces they add water back to the site at which they cut, called enzymatic hydrolysis, for proteins it is called proteolysis. The initial products of this hydrolysis are polypeptides, and smaller products are called simply peptides; these are called wheat protein hydrolysates. These hydrolysates can create allergens of wheat proteins that previously did not exist by the exposure of buried antigenic sites in the proteins.

When proteins are cut into polypeptides, buried regions are exposed to the surface, and these buried regions may possibly be antigenic. Such hydrolyzed wheat protein is used as an additive in foods and cosmetics. The peptides are often 1 kDa in size (9 amino acid residues in length) and may increase the allergic response (Akiyama et al., 2006). These wheat polypeptides can cause immediate contact urticaria in susceptible people (Laurière et al., 2006).

## **COMMON DISEASES**

### **Celiac disease**

The occurrence of celiac disease (CD) is about 1: 100 (Europe, USA) and higher than supposed earlier (Jankowiak et al., 2008). If we look back, the history of celiac disease can be separated into many ages, each driven by diagnostic advancement and deeper

knowledge of disease pathogenesis. These advances are paralleled by the recognition of new clinical patterns linked with CD and by an incessant redefinition of the pervasiveness of the disease in population. Due to its exclusive pathogenesis, celiac disease is extensively examined as a model immuno-genetic disorder (Mamone et al., 2011). Coeliac disease is believed to have a genetic component because there is a strong association of the disease with the presence of specific HLA class II antigens, predominantly HLA DQ2 and (to a lesser extent) DQ8 (Louka and Sollid, 2003). It is believed that the proteins implicated in the disease, the prolamins, bind to the DQ2 or DQ8 molecule, giving rise to T-cell activation and triggering a sequence of events leading to full-blown disease. It was found by Sjostrom et al. (1998) that T-cell reactivity is enhanced by the action of tissue transglutaminase, which deamidates specific glutamine residues of gliadin and glutenin peptides.

In the beginning, CD was believed as a chronic indigestion, even if the causal food was not identified. Later, the disease was confirmed to depend on intolerance to wheat gliadin that leads to typical mucosal changes in the gut and to a malabsorption syndrome. These facts directed to cure the disease with a gluten-free diet. After the recognition of antibodies to gluten in the serum of patients and the recognition of gluten-specific lymphocytes in the mucosa, CD was depicted as an autoimmune disorder that usually presents as enteropathy in genetically susceptible individuals like a chronic "gluten infection" (Zhernakova et al., 2010; Tommasini et al., 2011). It is

distinguished by polygenic predisposition, autoimmune nature, principally asymptomatic or atypical clinical course (Pavlović et al., 2010). CD4(+) T cells have a vital function in CD, identifying and binding complexes bearing gluten peptides that have endured digestion and are deamidated by tissue transglutaminase (TG2) thus supporting disease progression (Lindfors et al., 2010; Huan et al., 2011). The intestinal mucosa of susceptible individuals is crossed by undigested gluten that direct to a marked inflammatory response (Bas et al., 2009). One of the most serious side effects is osteopenia, while increased risk of malignant diseases (James and Scott, 2001; Pengiran Tengah et al., 2002; Vestergaard, 2003). As there are indications that a gluten-free diet protects coeliac patients against the development of these complications, the importance of compliance to such a diet is stressed. As there is no cure, a lifelong compliance to a gluten-free diet is the only way to alleviate symptoms. The innate immune system consists of a variety of immune responses that constitute the first line of defense against the invasion of foreign substances, e.g. pathogens. Its response to challenge is ahead of the induction of the adaptive immune response, is not antigen-specific and is without memory. Activation at inappropriate sites or to an excessive degree can exacerbate tissue damage in several diseases, and a possible role in the first stages of celiac disease has been suggested (Maiuri et al., 2003).

Typical celiac disease, distinguished by mild to harsh gastrointestinal symptoms, is less widespread than non-classic celiac disease, illustrated by absence of gastrointestinal

symptoms (Polanco, 2008; Snyder et al., 2008; Cabrera-Chávez et al., 2009; Silva et al., 2010). Coeliac disease consequences from composite interactions between the innate immune system, an adaptive T and B cell response and the mucosal barrier where inflammation is eventually manifested (Mothes et al, 2009; Heap et al., 2009; De Vincenzi et al., 2010).

Good devotion to gluten-free diet is compulsory to sustain the optimal bone health (Mora, 2008). A key mechanism of the disease is immune-mediated T-lymphocyte activation in the gastrointestinal mucosa following gluten ingestion. Inflammation of the intestinal mucosa and atrophy of the villi occur, and can result in the inadequate absorption of nutrients (Fasano and Catassi, 2001). Briefly, mechanism of causing allergy at bowel level includes:

- Glutaminase (an intestinal enzyme) deficiency
- Increased permeability of the bowel to macromolecules, including the allergen
- The production of antibodies to the relevant prolamins, or a fragment of it
- Increased production of inflammatory mediators

The onset of noticeable symptoms of CD seems to be dependent on the following: exposure to wheat, as when an infant is weaned (introduction of solids); predisposition through family history; and some kind of "trigger" mechanism (Fasano and Catassi, 2001; Palosuo et al. 2001). The clinical manifestations of the disease vary markedly with

the age of the affected person, the duration and severity of the disease and the presence of extra-intestinal pathologic conditions. The symptoms of CD include diarrhoea, bloating, weight loss, anemia, chronic fatigue, bone pain, and muscle cramps. Symptoms vary with respect to age from young to elder children (Fasano and Catassi, 2001). Corn, rice and other cereal grains such as sorghum, millet, teff, ragi and Job's tears as well as buckwheat, quinoa and amaranth can safely be ingested by a person with celiac disease (CD). Spelt and kamut however, should be avoided in CD (Thompson, 2000).

## **Baker's asthma**

Baker's asthma is an allergy caused by inhalation of cereal flour and is most common disease among workers with occupational exposure to cereal flour. Almost 4-25% of bakery workers are affected by prolonged exposure to wheat products. IgE-mediated reactions to wheat have been demonstrated for baker's asthma as early as the beginning of the 20th century (Houba, 1998). The insoluble gluten proteins are mainly responsible for these allergenic responses (Wiszniewska et al., 2011; Palosuo, 2003). Contact with or inhalation of wheat flour proteins is one of the causes of baker's asthma (an occupational allergy), but allergens other than the wheat itself may also be reasons. Symptoms of this disease include rhinitis, skin itching/rash, ocular symptoms (including tearing, itching and conjunctival injection), respiratory symptoms (including

coughing, wheezing, and shortness of breath and sputum production) and "grain fever" (Matsumura et al., 1994).

Hypersensitivity reactions have been noted in case of wheat flour, both by inhalation and ingestion (Merget et al., 2011; Baldo et al., 1980). Most prominent allergens for baker's asthma have been related to the salt-soluble fraction of wheat. Gomez *et al.* (1990) identified and characterized several 12-16 kDa salt-soluble proteins, members of the  $\alpha$ -amylase/trypsin inhibitor family, as major allergens associated with baker's asthma. In bakers' asthma, water-soluble proteins i.e. members of the  $\alpha$ -amylase inhibitor family, acyl-CoA oxidase, fructose-bisphosphatase aldolase (Weiss et al., 1997) and peroxidase (Sanchez-Monge et al., 1997) have been identified as important wheat allergens

## **Wheat-dependent exercise-induced reactions**

Wheat-dependent exercise-induced anaphylaxis is a separate form of wheat-allergy, which is induced by physical exercise after ingestion of wheat. Exercise within 3 hours of wheat consumption can induce an adverse reaction in susceptible individuals. In some cases, this can also occur when wheat is consumed directly after exercise. Typical symptoms experienced include asthma, urticaria, angioedema, dyspnoea, syncope and anaphylaxis (Palosuo et al., 2001).

In a previous study, Ioachim et al. (2002) demonstrated that the circulating levels of gliadin, which is recognized as an allergen for a variety of wheat allergies, were correlated with the clinical symptoms in patients with wheat-dependent exercise-induced anaphylaxis (WDEIA). Interestingly, increased serum gliadin levels induced by exercise and aspirin were also observed in healthy subjects. It is also hypothesized that aspirin facilitates the absorption of intact gliadin due to an increase in gastrointestinal permeability (Jones and Jones, 2000; Tsunoda et al., 2003). The typical symptom is generalized urticaria associated with shock or hypotension. Although the mechanism of physical exercise-induction in the anaphylaxis is indistinguishable, an immediate-type hypersensitivity to water/salt-insoluble fraction of wheat proteins (gluten) has been considered to underlie in this disease.

## **Atopic dermatitis**

Atopic dermatitis (AD) is a relapsing inflammatory skin disease characterized by eczematous skin lesions and intense pruritus (Patriarca *et al.*, 2007). In AD, pruritus is often resistant to H1-antihistamine treatment. Therefore other mediators, including neuropeptides, neurotransmitters, proteinases, arachidonic derivatives, and cytokines, have been suggested for a pathophysiologic role of pruritus in AD (Meglio et al., 2004).

Children with atopic disorders tend to have a higher prevalence of food allergy; about 35% of children with moderate to severe atopic dermatitis have IgE mediated food

allergy and about 6–8 % of asthmatic children have food-induced wheezing (Novembre et al., 1988) but 80% of them outgrow their food allergy. Later, Ellman et al. (2002) found that children with atopic dermatitis for a period of 10 years did not manifest reactions to an increasing variety of foods.

At the start of present decade, Wantanabe et al. (2000) reported the safety and usefulness of hypoallergenic wheat product in patient with atopic dermatitis (AD) having wheat allergy (Yamamoto et al., 2004)

## Conclusion

Wheat is consumed all over the world, considering as a staple food in many parts of the world. However, many people are intolerant to wheat proteins especially gluten protein. IgE-mediated wheat allergy is developed in the absence of oral tolerance. As a result of these reactions, different symptoms of allergy take place. Consequently, these symptoms result in different diseases like atopic dermatitis, celiac disease and baker's asthma. Baker's asthma is due to prolonged exposure of workers to the baking plant. Though, these diseases can be controlled by following different remedies. The preventive measures to avoid wheat allergy include consumption of fruits and vegetables and gluten free products. Additionally, avoid fried foods, sauces and all baked goods. Considering the nutshell of this review, various proteins present in wheat causes allergenic reaction in the community. So, the people who are intolerant to wheat

# ACCEPTED MANUSCRIPT

should avoid the proteins mentioned in this article. Though, further research is required to overcome the allergenic reactions caused by wheat as it is used as a staple food in many parts of the world and people cannot completely remove it from their diet.

## Literature Cited

Ajala, A.R., Cruz, A.G., Faria, J.A.F., Walter, E.H.M., Granato, D., Sant-Ana, A.S. (2010). Food allergens: Knowledge and practices of food handlers in restaurants. *Food Control*. **21**:1318–1321.

Akagawa, M., Handoyo, T., Ishii, T., Kumazawa, S., Morita, N., and Suyama, K. (2007). Proteomic analysis of wheat flour allergens. *J Agric Food Chem*. **55**:6863–6870.

Akiyama, M., Titeux, M., Sakai, K., McMillan, J.R., Tonasso, L., Calvas, P., Jossic, F., Hovnanian, A., and Shimizu, H. (2007). DNA-Based Prenatal Diagnosis of Harlequin Ichthyosis and Characterization of ABCA12 Mutation Consequences. *J Investigative Dermatology*. **127**:568-573.

Andera, A.E., Mannetje, A.T., Douwes, J., Cheng, S., McLean, D., Ellison-Loschmann, L., and Pearce N. (2009). The New Zealand workforce survey II: occupational risk factors for asthma. *Oxford J*. **54**(2):154-164.

[Ann Dermatol Venereol](#). 2010 Apr; **137**(4):281-4.

Anna, N.W., and Antonella, M. (2009). Food protein-induced enterocolitis syndrome. *Current Opinion in Allergy and Clinical Immunology*: **9**(4):371-377.

Armentia, A., Sanchez-Monge, R., Gomez, L., Barber, D., and Salcedo, G. (1993). *In vivo* allergenic activities of eleven purified members of a major allergen family from wheat and barley flour. *Clin Exp Allergy*. **23**(5):410–5

Atchison, J., Head, L., and Gates, A. (2010). Wheat as food, wheat as industrial substance; comparative geographies of transformation and mobility. *Geoforum*. **41**:236–246.

Baldo, B.A., Krilis, S., and Wrigley, C.W. (1980). Hypersensitivity to inhaled flour allergens. *Allergy*. **35**:45-56.

Bas, A., Forsberg, G., Sjöberg, V., Hammarström, S., Hernell, O., Hammarström, M.L. (2009). Aberrant extrathymic T cell receptor gene rearrangement in the small intestinal mucosa: a risk factor for coeliac disease? *Gut*. **58**(2):189-195.

Battais, F., Courcouxb, P.,ves Popineaua, Y., Moneret-Vautrinc, K.D.A., and Denery-Papini, S. (2005). Food allergy to wheat: differences in immunoglobulin E-binding proteins as a function of age or symptoms. *J Cereal Sci*. **42**:109–117.

Biagini, R.E., MacKenzie, B.A. Sammons, D.L., Smith, J.P., Striley, C.A., Robertson, S.K., and Snawder, J.E. (2004). Evaluation of the prevalence of anti- wheat anti-flour dust, and anti-amylase specific IgE antibodies in US blood donors. *Ann. Allergy Asthma Immunol*. **92**:649–653.

Bittner, C., Grassau, B., Frenzel, K., and Baur, X. (2010). Identification of wheat gliadins as an allergen family related to baker's asthma. *J Allergy Clinical Immunol.* **121**(3).

Blands, J., Diamant, B., Kallós, P., Kallós-Deffner, L., and Lowenstein, H. (1976). Flour allergy in bakers. I. Identification of allergenic fractions in flour and comparison of diagnostic methods. *Int Arch Allergy Appl Immunol.* **52**(1-4): 392–406.

Bouchez-Mahiout, I., Snégaroff, J., Tylichova, M., Pecquet, C., Branlard, G., and Laurière, M. (2010). Low Molecular Weight Glutenins in Wheat-Dependant, Exercise-Induced Anaphylaxis: Allergenicity and Antigenic Relationships with Omega 5-Gliadins. *Int Arch Allergy Immunol.* **153**:35-45.

Brant, A. (2007). Baker's asthma. *Curr. Opin. Allergy Clin Immunol.* **7**:152-155.

Branum, A., and Lukacs, S., (2008). Food allergy among U.S. children: Trends in prevalence and hospitalizations. NCHS Data Brief, No. 10. National Center for Health Statistics, Hyattsville, MD.

Bucka, J, Hattersley, S., and Kimber, I. (2010). Food allergy – science and policy needs – The UK Food Standards Agency. Research Programme. *Toxicol.* **278**:319-325.

Cabrera-Chávez, F., and de-Barca, A.M. (2009). Bovine milk intolerance in celiac disease is related to IgA reactivity to alpha- and beta-caseins. *Nutrition.* **25**(6):715-6.

Case, S. (2005). The gluten free diet: How to provide effective education and resources.

*Gastroenterol.* **128**(4):128-134.

Castellanos-Rubio, A., Santin, I., Irastorza, I., Castaño, L., Vitoria, J.C., Bilbao, J.R. (2009).

TH17 (and TH1) signatures of intestinal biopsies of CD patients in response to gliadin.

*Autoimmunity.* **42**:69–73.

Catassi and Fasano, 2008 C. Catassi and A. Fasano, Celiac disease. In: E.K. Arendt and F. Dal

Bello, Editors, Gluten-free cereal products and beverages, Academic Press, London

(2008).

Čelakovská, J., Vaněčková, J., Ettlrová, K., Ettler, K., and Bukač, J. (2010). The role of

atopy patch test in diagnosis of food allergy in atopic eczema/dermatitis syndrom

in patients over 14 years of age. *Acta Medica.* **53**(2):101–108.

Cho, H.N., Hong, S., Lee, S.H. and Yum, H.Y. 2011. Nutritional status according to sensitized food

allergens in children with atopic dermatitis. *Allergy Asthma Immunol. Res.* **3**(1): 53-7.

Ciclitira, P.J., Ellis, H.J., Lundin, K.E.A. (2005). Gluten free diet- what is toxic? *Best Practice Res*

*Clin Gastroenterol.* **19**(3):359-371.

Codex Alimentarius Commission 2007. Report of the 29th session of the Codex Committee on

Nutrition and Foods for Special Dietary uses. Draft revised Codex Standard for Foods

for Special Dietary Use for Persons Intolerant to Gluten. Codex Alimentarius Alinorm 08/31/26, para. 64 and Appendix III, Rome, Italy, pp. 50–51.

Codex Alimentarius Commission 2008. Report of the 31st session of the Codex Committee on Nutrition and Foods for Special Dietary uses. Revised Codex Standard for Foods for Special Dietary Use for Persons Intolerant to Gluten. Codex Alimentarius Alinorm 08/31/REP, Appendix VII, Geneva, Switzerland, p. 107.

Constantin, C., Quirce, S., Grote, M., Touraev, A., Swoboda, I., Stoecklinger, A., Mari, A., Thalhamer, J., Heberle-Bors, E., and Valenta, R. (2008). Molecular and Immunological Characterization of a Wheat Serine Proteinase Inhibitor as a Novel Allergen in Baker's Asthma. *J Immunol.* **180**:7451-7460.

Constantin, C., Quirce, S., Poorafshar, M., Touraev, A., Niggemann, B., Mari, A., Ebner, C., Akerström, H., Heberle-Bors, E., Nystrand, M.m and Valenta, R. (2009). Micro-arrayed wheat seed and grass pollen allergens for component-resolved diagnosis. *Allergy.* **64**:1030–1037.

Cosnes, J., Cellier, C., Viola, S., Colombel, J.F., Michaud, L., Sarles, J., Hugot, J.P., Ginies, J.L., Dabadie, A., Mouterde, O., Allez, M., Nion-Larmurier, I., and The Groupe D'Etude et de Recherche Sur la Maladie Coeliaque. (2008). Incidence of autoimmune disease in celiac disease: Protective effect of the gluten free diet. *Clinical Gastroenterol Hepatol.* **6**(7):753-758.

- Cross, A.H., and Golumbek, P.T. (2003). Neurologic manifestations of celiac disease Proven, or just a gut feeling? *Neurol.* **60**:1566–1568.
- De-Benedetto, A., Agnihotri, R., McGirt, L.Y., Bankova, L.G., and Beck, L.A. (2009). Atopic dermatitis: a disease caused by innate immune defects. *J Invest Dermatol.* **129**:14-30.
- Denery-Papini, S., Laurière, M., Branlard, G., Morisset, M., Pecquet, C., Choudat, D., Merlino, M., Pineau, F., Popineau, Y., Boulenc, E., Bouchez-Mahiout, I., Bodinier, M., and Moneret-Vautrin, D.A. (2007). Influence of the allelic variants encoded at the Gli-B1 locus, responsible for a major allergen of wheat, on IgE reactivity for patients suffering from food allergy to wheat. *J. Agric Food Chem.* **55**(3):799–805.
- De-Vincenzi, M., Vincentini, O., Di-Nardo, G., Boirivant, M., Gazza, L., and Pogna, N. (2010). Two prolamin peptides from durum wheat preclude celiac disease-specific T cell activation by gluten proteins. *Eur J Nutr.* **49**(4):251-5.
- Dewar, D.H., Amato, M., Ellis, J., Pollock, E.L., Gonzalez-Cinca, N., Wieser, H., and Ciclitira, P.J. (2006). The toxicity of high molecular weight glutenin subunits of wheat to patients with coeliac disease. *EuropJ Gastroenterol Hepatol.* **18**:483–491.
- Dickerson, F., Stallings, C., Origoni, A., Vaughan, C., Khushalani, S., Leister, F., Yang, S., Krivogorsky, B., Alaedini, A., and Yolke, R. (2010). Markers of Gluten Sensitivity and Celiac Disease in Recent-Onset Psychosis and Multi-Episode Schizophrenia. *Biol psychiatry.* **68**:100–104.

Dieterich, W., Esslinger, B., Trapp, D., Hahn, E., Huff, T., Seilmeier, W., Wieser, H., and Schuppan, D. (2006). Cross linking to tissue transglutaminase and collagen favours gliadin toxicity in coeliac disease. *Gut*. **55** (4): 478–484.

Dong, P., Reed, S. A., Yi, S.A., Kalmykov, S., Shvets, G., and Downer, M.C. (2010). Formation of optical bullets in laser-driven plasma bubble accelerators. *Phys Rev Lett*. **104**(13):134801.

Donovan, G.R., and Baldo, B.A. (1990). Crossreactivity of IgE antibodies from sera of subjects allergic to both ryegrass pollen and wheat endosperm proteins: evidence for common allergenic determinants. *Clin Exp Allergy*. **20**(5):501–9.

Droste, J., Myny, K., Van Sprundel, M., Kusters, E., Bulat, P., Braeckman, L., Vermeire, P., and Vanhoorne, M. (2003). Allergic sensitization, symptoms, and lung function among bakery workers as compared with a nonexposed work population. *J Occup Environ Med*. **45**:648–655.

Dykewicz, M.S., and Hamilos, D.L. (2010). Rhinitis and sinusitis. *J Allergy Clin Immunol*. **125**(2):103–115.

Elder, J.H. (2008). "The gluten-free, casein-free diet in autism: an overview with clinical implications". *Nutr Clin Pract* **23** (6): 583–538.

Epub 2010 Feb 18. [Contact urticaria induced by hydrolyzed wheat proteins in cosmetics].

[Olaiwan A](#), [Pecquet C](#), [Mathelier-Fusade P](#), [Francès C](#).

Esch, R.E. (2004). Grass pollen allergens. *Clin Allergy Immunol* . **18**:185-205.

- Fernandez-Jimenez, N., Castellanos-Rubio, A., Plaza-Izurietta, L., Gutierrez, G., Castano, L., Vitoria, J.C., and Bilbao, J.R. (2010). Analysis of  $\beta$ -defensin and Toll-like receptor gene copy number variation in celiac disease. *Human Immunol.* **71**: 833–83.
- Gao, L., Wang, A., Li, X., Dong, K., Wang, K., Appels, R., Ma, W., and Yan, Y. (2009). Wheat quality related differential expressions of albumins and globulins revealed by two-dimensional difference gel electrophoresis (2-D DIGE). *J Proteomics.* **73**(2):279-296.
- Grant, E.C. (1979). "Food allergies and migraine". *Lancet* 1 (**8123**): 966–969.
- Hadjivassiliou, M., Grünewald, R.A., Kandler, R.H., Chattopadhyay, A.K., Jarratt, J.A., Sanders, D.S., Sharrack, B., Wharton, S.B., and Davies-Jones, G.A.B. (2006). Neuropathy associated with gluten sensitivity. *J Neurol Neurosurg Psychiatry.* **77**:1262-1266.
- Hadjivassiliou, M., Rao, D.G., Wharton, S.B., Sanders, D.S., Grünewald, R.A., and Davies-Jones, A.G.B. (2010). Sensory ganglionopathy due to gluten sensitivity. *Neurol.* **75**:1003-1008.
- Hansen, I., Klimek, L., Mosges, R., and Hormann, K. (2004). Mediators of inflammation in the early and the late phase of allergic rhinitis. *Curr Opin Allergy Clin Immunol.* **4**(3):159-63.
- Heap, G.A., and Van-Heel, D.A. (2009). Genetics and pathogenesis of coeliac disease. *Semin Immunol.* **21**(6):346-54.

- Heffler, E., Nebiolo, F., Asero, R., Guida, G., Badiu, I., Pizzimenti, S., Marchese, C., Amato, S., Mistrello, G., Canaletti, F. and Rolla, G. 2011. Clinical manifestations, co-sensitizations, and immunoblotting profiles of buckwheat-allergic patients. *Allergy*. **66**(2): 264-70.
- Helm, R., and Burks, A. (2000). Mechanisms of food allergy. *Curr Opin. Immunol*. **12**: 647–653.
- Heratizadeh, A., Wichmann, K. and Werfel, T. 2011. Food allergy and atopic dermatitis: how are they connected? *Curr. Allergy Asthma Rep*. **11**(4): 284-91.
- Holtmeier, W., and Caspary, W.F. (2006). Celiac disease. *Orphanet J Rare Dis*. **1**:3. Huan, J., Meza-Romero, R., Mooney, J.L., Vandenbark, A.A., Offner, H., Burrows, G.G. (2011). Single-chain recombinant HLA-DQ2.5/peptide molecules block  $\alpha$ 2-gliadin-specific pathogenic CD4+ T-cell proliferation and attenuate production of inflammatory cytokines: a potential therapy for celiac disease. *Mucosal Immunol*. **4**(1):112-20.
- Hvatum, M., Kanerud, L., Hällgren, R., and Brandtzaeg, P. (2006). The gut-joint axis: cross reactive food antibodies in rheumatoid arthritis. *Gut*. **55**(9):1240–7.
- Imai, T. and Y. Likura. (2003). The national survey of immediate type of food allergy. *Arerugi*. **52**:1006-1013.
- Isolauri, E., Kalliomaki, M., Rautava, S., Salminen, S., and Laitinen, K. (2009). Obesity: extending the hygiene hypothesis. *Nestle Nutr Workshop Ser*. **64**:75–89.

- Jabri, B., Sollid, L.M. (2009). Tissue-mediated control of immunopathology in celiac disease. *Nat Rev Immunol.* **9**:858–70.
- Jankowiak, C., Ludwig, D. (2008). Frequent causes of diarrhea: celiac disease and lactose intolerance. *Med Klin (Munich).* **103**(6):413-22.
- Jones, S.M., Magnolfi, C.F., Cooke, S.K., and Sampson, H.A. (1995). Immunologic cross-reactivity among cereal grains and grasses in children with food hypersensitivity. *J Allergy Clin Immunol.* **96**:341–51.
- Karatay, S., Erdem, T., Kiziltunc, A., Mugnaini, L., Caffarelli, C., Cavagni, G., Giovane, A., and Vierucci, A. (2006). General or personal diet: the individualized model for diet challenges in patients with rheumatoid arthritis. *Rheumatol. Int.* **26**(6):556–60.
- Kasarda, D.D., Autran, J., Lew, E.J.L., Nimmo, C.C., and Shewry, P.R. (1983). N-terminal amino acid sequences of u-gliadins and u-secalins. Implications for the evolution of prolamin genes. *Biochimica Biophysica Acta.* **747**:138-150.
- Keet, C.A., and Wood, R.A. (2009). Risk factors for peanut allergy. *J Allergy Clin Immunol.* **24**(2):387.
- Kusaba, M., Miyahara, K., Iida, S., Fukuoka, H., Takano, T., Sassa, H., Nishimura, M., and Nishio, T. (2003). Low glutelin content1: a dominant mutation that suppresses the glutelin multigene family via RNA silencing in rice. *Plant Cell.* **15**:1455-1467.

Lack, G. (2008). Epidemiologic risks for food allergy. *J Allergy Clin Immunol.* **121**:1331-1336.

Langeland, T., and Nyruud, M., (1982). Contact urticaria to wheat bran bath: a case report. *Acta Derm. Venereol.* **62** (1):82-3.

Laurière, M., Pecquet, C., Bouchez-Mahiout, I., Snégaroff, J., Bayrou, O., Raison-Peyron, N., and Vigan, M. (2006). Hydrolysed wheat proteins present in cosmetics can induce immediate hypersensitivities. *Contact Derm.* **54**(5):283-9.

Lehto, M., Airaksinen, L., Puustinen, A., Tillander, S., Hannula, S., Nyman, T., Toskala, E., Alenius, H., and Lauerma, A. (2010). Thaumatin-like protein and baker's respiratory allergy. *Annals of Allergy Asthma Immunol.* **104**(2):139-146.

Leon, F. (2011). Flow cytometry of intestinal intraepithelial lymphocytes in celiac disease. *J Immunol Methods.* 363:177-187.

Lindfors, K., Mäki, M., and Kaukinen, K. (2010). Transglutaminase 2-targeted autoantibodies in celiac disease: Pathogenetic players in addition to diagnostic tools? *Autoimmun Rev.* **9**(11):744-9.

Loussert, C., Popineau, Y., and Mangavel, C. (2008). Protein bodies ontogeny and localization of prolamin components in the developing endosperm of wheat caryopses. *J Cereal Sci.* **47**:445-456.

Maiuri, L., Ciacci, C., Ricciardelli, I., Vacca, L., Raia, V., Auricchio, S., Picard, J., Osman, M., Quarantino, S., and Londei, M. (2003). Association between innate response to gliadin and activation of pathogenic T cells in coeliac disease. *Lancet*. **362**:30–7.

Majithia, V., and Geraci, S.A. (2007). Rheumatoid arthritis: diagnosis and management. *Am. J. Med.* **120**(11):936–939.

Mamone, G., Picariello, G., Addeo, F., and Ferranti, P. (2011). Proteomic analysis in allergy and intolerance to wheat products. *Expert Rev Proteomics*. **8**(1):95-115.

Mangiavillano, B., Masci, E., Parma, B., Barera, G., Viaggi, P., Albarello, L., Tronconi, G.M., Mariani, A., Testoni, S., Santoro, T., and Testoni, P.A. (2010). Bulb biopsies for the diagnosis of celiac disease in pediatric patients. *Gastrointestinal Endoscopy*. **72**(3):564-568.

Matsuo, H., Morimoto, K., Akaki, T., Kaneko, S., Kusatake, K., Kuroda, T., Niihara, H., Hide, M., and Morita, E. (2005). Exercise and aspirin increase levels of circulating gliadin peptides in patients with wheat-dependent exercise-induced anaphylaxis. *Clin Exp Allergy*. **35**(4):461-466.

Matsuo, H., Morita, E., Tatham, A.S., Morimoto, K., Horikawa, T., Osuna, H., Ikezawa, Z., Kaneko, S., Kohno, K., and Dekio, S. (2004). "Identification of the IgE-binding

epitope in omega-5 gliadin, a major allergen in wheat-dependent exercise-induced anaphylaxis. *J Biol Chem.* **279**(13):12135–12140.

May. J.R., and Smith, P.H. Allergic Rhinitis. In: Dipiro, J.T., Talbert, R.L., Yee, G.C., et al eds. *Pharmacotherapy: A Pathophysiologic Approach*. 7th ed. New York, NY: The McGraw-Hill Companies, Inc **2008**:1565-1575.

Merget, R., Sander, I., van-Kampen, V., Beckmann, U., Heinze, E., Raulf-Heimsoth, M. and Bruening, T. 2011. Allergic asthma after flour inhalation in subjects without occupational exposure to flours: an experimental pilot study. *Int. Arch. Occup. Environ. Health* **84**(7): 753-60.

Metz-Favre, C., Linhart, B., Focke-Tejkl, M., Purohit, A., de Blay, F., Valenta, F., and Pauli, G. (2007). Skin test diagnosis of grass pollen allergy with a recombinant hybrid molecule. *J Allergy Clin Immunol.* **120**(2):315-321.

Mittag, D., Niggemann, B., Sander, I., Reese, I., Fiedler, E.M., Worm, M., Vieths, S., and Reese, G. (2004). "Immunoglobulin E-reactivity of wheat-allergic subjects (baker's asthma, food allergy, wheat-dependent, exercise-induced anaphylaxis) to wheat protein fractions with different solubility and digestibility. *Mol Nutr Food Res.* **48** (5):380–389.

Moodie, S.J., Norman, P.J., King, A.L., Fraser, J.S., Curtis, D., Ellis, H.J., Vaughan, R.W., and Ciclitira, P.J. (2002). Analysis of candidate genes on chromosome 19 in coeliac disease: An association study of the KIR and LILR gene clusters. *Eur J Immunogenet.* **29**:287-91

Mora, S. (2008). Celiac disease in children: impact on bone health. *Rev Endocr Metab Disord.* **9**(2):123-130.

Morita, E., Kunie, K., and Matsuo, H. (2007). Food-dependent exercise-induced anaphylaxis. *J Dermatol Sci.* **47**(2):109-117.

Mothes, T., Uhlig, H.H., and Richter, T. (2009). Recent aspects of antibody determination for the diagnosis of coeliac disease. *Dtsch Med Wochenschr.* **134**(30):1525-8.

Mulitz, R.M., Pratt, D.S., and Schocket, A.L. (1979). Exercise-induced ana-phylactic reaction to shellfish. *J Allergy Clin Immunol.* **63**: 433-434.

Niggemann, B., Reibel, S., Roehr, C.C., Felger, D., Ziegert, M., Sommerfeld, C., and Wahn, U. (2001). Predictors of positive food challenge outcome in non-IgE-mediated reactions to food in children with atopic dermatitis. *J Allergy Clin Immunol.* **108**:1053-1058.

Novembre, E., Cianferoni, A., Bernardini, R., Mugnaini, L., Caffarelli, C., Cavagni, G., Giovane, A., and Vierucci, A. (1998). Anaphylaxis in Children: Clinical and Allergologic Features. *Pediatrics*. **101**(4).

Oita, S., Hayashi, T., and Ohnishi-Kameyama, M. (2009). Degradation of epitope peptides of wheat gliadin and glutenin for atopic dermatitis by crude proteases from germinated wheat seeds. *Food Sci Technol Res*. **15**:639-644.

Pacharn, P., Jirapongsananuruk, O., Daengsuwan, T., Vichyanond P., and Visitsunthorn, N. (2009). Wheat-dependent, exercise-induced anaphylaxis in Thai children: A report of 5 Cases. *Asian Pacific J Allergy Immunol*. **27**: 115-120.

Palacin, A., Quirce, S., Sánchez-Monge, R., Fernández-Nieto, M., Varela, J., Sastre, J., and Salcedo, G. (2008). Allergy to kiwi in patients with baker's asthma identification of potential cross-reactive allergens. *Annals of Allergy Asthma Immunol*. **101**:200-205.

Palacin, L., Prada, C.D., Syafiie, S., and Tadeo, F. (2008). Library for Dynamic Simulation of Reverse Osmosis Plants, Proc. of 20th European Modeling and Simulation Symposium. Sep. 17-19, Briatico, Italy, 2008.

Pascual, J., and Leno, C. (2005). A woman with daily headaches". *The journal of headache and pain: official journal of the Italian Society for the Study of Headaches*. **6**(2):91-92.

- Pastorello, E.A., Farioli, L., Conti, A., Pravettoni, V., Bonomi, S., Iametti, S., Fortunato, D., Scibilia, J., Bindslev-Jensen, C., Ballmer-Weber, B., Robino, A.M., and Ortolani, C. (2007). Wheat IgE-mediated food allergy in European patients: alpha-amylase inhibitors, lipid transfer proteins and low-molecular-weight glutenins. Allergenic molecules recognized by double-blind, placebo-controlled food challenge. *Int Arch Allergy Immunol.* **144**(1):10–22.
- Paula, R.A., George, D.T. (2010). Pathophysiology, diagnosis and management of exercise-induced anaphylaxis. *Current Opinion Allergy Clin Immunol.* **10**(4):312–317.
- Pavlović, M., Radlović, N., Leković, Z., Berenji, K., Stojsić, Z., Radlović, V. (2010). Coeliac disease as the cause of resistant sideropenic anaemia in children with Down's syndrome: case report. *Srp Arh Celok Lek.* **138**(1-2):91-4.
- Pérez-Pimiento, A.J., Rodríguez-Cabreros, M.I., Lombardero, M., Prieto, L., Reaño, M., and Iglesias, A. (2007). Late-Onset Food Hypersensitivity to Wheat Flour. *J Investig Allergol Clin Immunol.* **17**(3): 202-206.
- Pizzuti, D., Senzolo, M., Buda, A., Chiarelli, S., Giacomelli, L., Mazzon, E., Curioni, A., Faggian, D. and De-Lazzari F. 2011. In vitro model for IgE mediated food allergy. *Scand. J. Gastroenterol.* **46**(2): 177-87.
- Polanco, I. (2008). Celiac disease. *J Pediatr Gastro Nutr.* **47**(1):3-6.

- Ramesh, S. (2008). Food allergy overview in children. *Clin Rev Allergy Immunol.* **34**(2):217-230.
- Rodrigo, L., Rodrigo, (2006). Celiac disease. *World J Gastroenterol.* **12**(41): 6585–6593.
- Rona, R., Keil, T., Summers, C., Gislason, D., Zuidmeer, L., Sodergren, E., Sigurdardottir, S., Lindner, T., Goldhahn, K., Dahlstrom, J., McBride, D., and Madsen, C. (2007). The prevalence of food allergy: a meta-analysis. *J Allergy Clin Immunol.* **120**:638–646.
- Rose, C., Armbruster, F.P., Ruppert, J., Igl, B.W., Zillikens, D., and Shimanovich, I. (2009). Autoantibodies against epidermal transglutaminase are a sensitive diagnostic marker in patients with dermatitis herpetiformis on a normal or gluten free diet. *J Am Academy Dermatol.* **61**(1): 39-43.
- Sampson, M.A., Munoz-Furlong, A., and Sicherer, S.H., (2006). Risk-taking and coping strategies of adolescents and young adults with food allergy. *J Allergy Clin Immunol.* **117**:1440–1445.
- Santin, I., Castellanos-Rubio, A., Perez deNanclares, G., Vitoria, J.C., Castaño, L., and Bilbao, J.R. (2007). Association of KIR2DL5B gene with celiac disease supports the susceptibility locus on 19q13.4. *Genes Immun.* **8**:171–176.
- Santin, I., de Nanclares, G.P., Calvo, B., Gaafar, A., Castaño, L., GEPV-N Group, Bilbao, J.R. Killer cell immunoglobulin-like receptor (KIR) genes in the Basque population:

Association study of KIR gene contents with type 1 diabetes mellitus. *Hum Immunol.* **67**:118–24.

Scibilia, J., Pastorello, E.A., Zisa, G., Ottolenghi, A., Ballmer-Weber, B., Pravettoni, V., Scovena, E., Robino, A., and Ortolani, C. (2008). Maize food allergy: a double-blind placebo-controlled study. *Clin Experi Allergy.* **38**:1943–1949.

Scibilia, J., Pastorello, E.A., Zisa, G., Ottolenghi, A., Bindslev-Jensen, C., Pravettoni, V., Scovena, E., Robino, A., and Ortolani, C. (2006). Wheat allergy: A double-blind, placebo-controlled study in adults. *J Allergy Clin Immunol.* **117**(2):433-439.

Shang, A., Huwiler-Müntener, K., Nartey, L., Jüni, P., Dörig, S., Sterne, J.A., Pewsner, D., and Egger, M. (2005). Are the clinical effects of homoeopathy placebo effects? Comparative study of placebo-controlled trials of homoeopathy and allopathy. *Lancet.* **366**(9487):726-732.

Shewry, P.R., (1995). Plant storage proteins. *Biol Rev Camb Philos Soc.* **70**:375–426.

Shewry, P.R., and Halford, N.G. (2002). Cereal grain storage proteins: structures, properties and role in grain utilization. *J Experimental Botany.* **53**:947–958.

Shewry, P.R., Tatham, A.S., Forde, J., Kreis, M., and Mifflin, B.J. (1986). The classification and nomenclature of wheat gluten proteins: a reassessment. *J Cereal Sci.* **4**:97-106.

Sicherer, S.H., Morrow, E.H., and Sampson, H.A. (2000). Dose-response in double-blind, placebo-controlled oral food challenges in children with atopic dermatitis. *J Allergy Clin Immunol.* **105**:582-586.

Sicherer, S.H., Sampson, H.A., 2010. Food allergy. *Allergy Clin. Immunol.* **125**, S116–S125

Sih, T., and Mion, O. (2010). Allergic rhinitis in the child and associated comorbidities. *Pediatric Allergy Immunol.* **21**(1):107-113.

Silva, T.S., and Furlanetto, T.W. (2010). Diagnosis of celiac disease in adults. *Rev Assoc Med Bras.* **56**(1):122-6.

Singh, J., Blundell, M., Tanner, G., and John, H. S. (2001). Albumin and Globulin Proteins of Wheat Flour: Immunological and N-terminal Sequence Characterisation. *J Cereal Sci.* **34**: 85–103

Smith, T.A. (2004). Preventing baker's asthma: an alternative strategy. *Occup Med.* **54**:21–27.

Snyder, C.L., Young, D.O., Green, P.H.R., and Taylor, A.K. (2008). Celiac Disease. Gene Reviews, University of Washington, USA.

Sotkovsky, P., Sklenar, J., Halada, P., Cinova, J., Setinova, I., Kainarova, A., Golias, J., Pavlaskova, K., Honzova, S. and Tuckova, L. 2011. A new approach to the isolation and characterization of wheat flour allergens. *Clin. Exp. Allergy* **41**(7): 1031-43.

Sutton, R., Hill, D.J., Baldo, B.A., Wrigley, C.W. (1982). Immunoglobulin E antibodies to ingested cereal flour components: studies with sera from subjects with asthma and eczema. *Clin Allergy*. **12**(1):63–74.

Tanaka, M., Nagano, T., Yano, H., Matsuda, T., Ikeda, T.M., Haruma, K. and Kato Y. 2011. Impact of  $\omega$ -5 gliadin on wheat-dependent exercise-induced anaphylaxis in mice. *Biosci. Biotechnol. Biochem.* **75**(2): 313-7.

Taylor, P.E., Flagan, R.C., Valenta, R., and Glovsky, M.M. (2002). Release of allergens as respirable aerosols: a link between grass pollen and asthma. *J Allergy Clin Immunol.* **109**:51-6.

Tommasini, A., Not, T., and Ventura, A. (2011). Ages of celiac disease: From changing environment to improved diagnostics. *World J Gastro.* **28**17(32):3665-71.

Tordesillas, L., Sirvent, S., Díaz-Perales, A., Villalba, M., Cuesta-Herranz, J., Rodríguez, R. and Salcedo, G. 2011. Plant Lipid Transfer Protein Allergens: No Cross-Reactivity between

Those from Foods and Olive and Parietaria Pollen. *Int. Arch. Allergy Immunol.* **56**(3): 291-296.

Valero Santiago, A., Amat Par, P., Sanosa Valls, J., Sierra Martínez, P., Malet Casajuana, A., and García Calderón, P.A. (1988). Hypersensitivity to wheat flour in bakers. *Allergologia et immunopathologia.* **16**(5): 309–14.

Valet, R.S., and Fahrenholz, J.M. (2009). Allergic rhinitis: update on diagnosis. *Consultant.* **49**:610-613

Van-der-Slik, A.R., Koeleman, B.P., Verduijn, W., Bruining, G.J., Roep, B.O., Giphart, M.J. (2003). KIR in type 1 diabetes: Disparate distribution of activating and inhibitory natural killer cell receptors in patients versus HLA-matched control subjects. *Diabetes.* **52**:2639-42.

Van-Eckert, R., Bond, J., Rawson, P., Klein, C., Stern, C., and Jordan, T.W. (2010). Reactivity of gluten detecting monoclonal antibodies to a gliadin reference material. *J Cereal Sci.* **51**(2):198-204.

Varjonen, E., Vainio, E., and Kalimo, K. (2000). Antigliadin IgE: indicator of wheat allergy in atopic dermatitis. *Allergy.* **55**:386-391.

- Walsh, B.J., Wrigley, C.W., Musk, A.W., and Baldo, B.A. (1985). A comparison of the binding of IgE in the sera of patients with bakers' asthma to soluble and insoluble wheat-grain proteins. *J Allergy Clin Immunol.* **76**(1):23–8.
- Wang, W., Han, J., Wu, Y., Yuan, F., Chen, Y. and Ge, Y. 2011. Simultaneous detection of eight food allergens using optical thin-film biosensor chips. *J. Agric. Food Chem.* **59**(13): 6889-94.
- Watanabe, M. (1993). Hypoallergenic rice as a physiologically functional food. *Trends in Food Sci Technol.* **4**(5):125-128.
- Wieser, H. (2007). Chemistry of gluten proteins. *Food Microbiol.* **24**(2):115-119.
- Wiszniewska, M., Nowakowska-Świrta, E., Pałczyński, C. and Walusiak-Skorupa, J. 2011. Diagnosing of bakers' respiratory allergy: is specific inhalation challenge test essential? *Allergy Asthma Proc.* **32**(2): 111-8.
- Wong, G.K.Y., Huissoon, A.P., Goddard, S., Collins, D.M., and Krishna, M.T. (2010). Wheat dependent exercise induced anaphylaxis: is this an appropriate terminology? *J Clin Pathol.* **63**:814-817.
- Yamamoto, M., Haruna, T., Yasui, K., Takahashi, H., Iduhara, M., Takaki, S., Deguchi, M., and Arimura, A. (2007). A novel atopic dermatitis model induced by topical

application with dermatophagoides farinae extract in NC/Nga mice. *Allergology International*. **56**:139–148.

Yazicioglu, M., Oner, N., Celtik, C., Okutan, O., and Pala, O. (2004). Sensitization to common allergens, especially pollens, among children with respiratory allergy in the Trakya region of Turkey. *Asian Pac. J. Allergy Immunol.* **22**(4):183–90.

Young, E., Stoneham, M.D., Petruckevitch, A., Barton, J., and Rona, R. (1994). A population study of food intolerance. *Lancet*. **343**:1127–1130.

Zeiger, R.S., Heller, S., Mellon, M.H., Forsythe, A.B., O'Connor, R.D., Hamburger, R.N., and Schatz, M. (1989). Effect of combined maternal and infant food-allergen avoidance on development of atopy in early infancy: a randomized study. *J. Allergy Clin. Immunol.* **84**(1):72–89.

Table 1: Percentages of prolamin type in different cereals

Cereals	Prolamin type	Amount (%)
Wheat	Gliadin	69
Corn	Zein	55
Barley	Hordein	46-52
Sorghum	Kafirin	52
Rye	Secalinin	30-50
Millet	Panicin	40
Oats	Avenin	16
Rice	Orzenin	5


Table 2: Monosaccharides present in eukaryotic glycoproteins

Sugar	Type	Abbreviation
<u><math>\beta</math>-D-Glucose</u>	<u>Hexose</u>	Glc
<u><math>\beta</math>-D-Galactose</u>	Hexose	Gal
<u><math>\beta</math>-D-Mannose</u>	Hexose	Man
<u><math>\alpha</math>-L-Fucose</u>	<u>Deoxyhexose</u>	Fuc
<u>N-Acetylgalactosamine</u>	<u>Aminohexose</u>	GalNAc
<u>N-Acetylglucosamine</u>	Aminohexose	GlcNAc
<u>N-Acetylneuraminic acid</u>	<u>Aminononulosonic acid</u> (Sialic acid)	NeuNAc
<u>Xylose</u>	<u>Pentose</u>	Xyl

CUTANEOUS EXPOSURE



Allergen



Skin-draining lymph nodes



ALLERGY

Th2 memory




Figure 1: Mechanism of wheat allergy

(Lack, 2008)